Experimental Study for Lithosphere-Atmosphere-Ionosphere Coupling : Observation of Atmospheric Parameters at Asahi Station, Chiba, Japan

*大村 潤平¹、韓 鵬²、吉野 千恵¹、服部 克巳¹、下 道國³、小西 敏春⁴、古屋 隆一⁵
*Junpei Omura¹, Peng Han², Chie Yoshino¹, Katsumi Hattori¹, Michikuni Shimo³, Toshiharu Konishi⁴, Ryuichi Furuya⁵

- 1. 千葉大学大学院理学研究科、2. 統計数理研究所、3. 藤田保健衛生大学、4. 応用光研工業株式会社、5. コムシステム株式会社
- 1. Graduate School of Science, Chiba University, 2. Institute of Statistical Mathematics, 3. Fujita Health University, 4. OHYO KOKEN KOGYO CO., LTD., 5. COM SYSTEM, INC.

The lonospheric anomaly is one of the most promising precursory phenomena for large earthquakes. Lithosphere-Atmosphere-Ionosphere Coupling (LAIC) model has been proposed to explain these phenomena. To examine the possibility of chemical channel of LAIC through the monitoring of atmospheric electricity parameters, we have installed sensors for the atmospheric electric field (AEF), atmospheric ion concentration (AIC), radon concentration, radon exhalation quantity (REQ), and weather elements. We will report the properties of variation in atmospheric electricity parameters observed at Asahi station (ASA), Japan to identify earthquake-related signals in these parameters. We found that the variation of radon exhalation quantity shows a clear negative correlation with 3 hours delay to the air pressure variation in clear days. Each season differs in daily pattern. AIC and AEF variations show lag correlation with radon exhalation quantity variation. To extract anomalous radon variation related to earthquakes, we should set a network of Radon monitoring and establish a model of radon variation for the future detailed analysis. We also observed cases that AEF has showed a spike-like

キーワード: Lithosphere-Atmosphere-Ionosphere Coupling、atmospheric ion concentration、atmospheric electric field、radon exhalation quantity

Keywords: Lithosphere-Atmosphere-Ionosphere Coupling, atmospheric ion concentration, atmospheric elctric field, radon exhalation quantity

increase at the same time as the time when AIC has largely increased. It must be going to be checked whether AEF data was taken in fair-weather period, however, it is suggested that change in local charge

distribution may have influenced AEF.