

Vertical motion of the neutral atmosphere above Tromsø

*野澤 悟徳¹、小川 泰信²、藤原 均³、川原 琢也⁴、津田 卓雄⁵、斎藤 徳人⁶、和田 智之⁶、高橋 透²、堤雅基²、川端 哲也¹、Hall Chris⁷、Brekke Asgeir⁷

*Satonori Nozawa¹, Yasunobu Ogawa², Hitoshi Fujiwara³, Takuya Kawahara⁴, Takuo T. Tsuda⁵, Norihito Saito⁶, Satoshi Wada⁶, Toru Takahashi², Masaki Tsutsumi², Tetsuya Kawabata¹, Chris Hall⁷, Asgeir Brekke⁷

1. 名古屋大学宇宙地球環境研究所、2. 国立極地研究所、3. 成蹊大学理工学部、4. 信州大学工学部、5. 電気通信大学情報理工学部、6. 理化学研究所光量子工学研究領域、7. UiT ノルウェー北極大学

1. Institute for Space-Earth Environmental Research, Nagoya University, 2. National Institute of Polar Research, 3. Faculty of Science and Technology, Seikei University, 4. Faculty of Engineering, Shinshu University, 5. The University of Electro-Communications, 6. RIKEN, 7. UiT The Arctic University of Norway

We will present results of vertical motion above Tromsø (69.6 deg. N, 19.2 deg. E) mainly based on sodium LIDAR data. Vertical motion of the neutral gases in the upper mesosphere and lower thermosphere (MLT) is a peculiar issue, and its understanding is important in terms of substance transport as well as thermal structures. Observations of the vertical wind in the MLT region are rather difficult, because vertical velocities are generally thought to be about two orders smaller than horizontal wind velocities. It is believed that the cold summer mesopause is set up by upward wind with strength of a few cm/s in the mesosphere. During high auroral activity intervals, some observations conducted by Fabry-Perot Interferometer (FPI) reported about 10 m/s or larger vertical wind velocity in the polar lower thermosphere. FPI measurements, however, suffer from a serious weakness of passive measurements: no information on the height observed. On the other hand, observations of vertical winds by radars are also difficult. Thus, our understanding of the vertical motion in the polar MLT region is still limited. The sodium LIDAR operated at Tromsø is capable of simultaneous measurements of wind velocities with five directions with a good accuracy (1-2 m/s). By using the LIDAR data (about 2100 hr data) obtained from October 2012 to March 2016 together with EISCAT, MF, and meteor radar data as well as auroral image data, we will discuss the characteristics of the vertical motion in the polar MLT.

We have found some events where the vertical wind blew with strength of about 10 m/s. In the case of January 14, 2015, the upward vertical wind with an amplitude of 10 m/s was found between 92 and 101 km over a few hours. During the night, the semidiurnal tide was strong with an amplitude of 100 m/s. This would confirm that strong vertical motion exists when such waves pass by the MLT region. In another event found in February 8, 2013, upward flows were observed between 94 and 96 km at the same time for 15 min, while no vertical flows were found at and above 97 km and at and below 93 km. Of particular interest in both cases is that a sporadic sodium layer (SSL) appeared nearby the height region where the upward vertical wind was observed at the same time (in the case of January 14, 2015) or 15 min later (in the case of February 8, 2013). In this presentation, we will address what conditions are needed for the vertical motion occurring, and also discuss possible relationship with the advent of SSLs.

キーワード：鉛直風、中間圏・下部熱圏、ライダー、トロムソ、EISCAT

Keywords: Vertical wind, Mesosphere and lower Thermosphere, LIDAR, Tromsø, EISCAT