

共役点オーロラにおける渦の発達 of 南北非対称性

North-south asymmetry of vortex evolution in conjugate aurora

*内田 ヘルベルト陽仁¹、片岡 龍峰^{2,1}、門倉 昭^{2,1}、小川 泰信^{2,1}、福田 陽子³、三好 由純⁴

*Herbert Akihito Uchida¹, Ryuho Kataoka^{2,1}, Akira Kadokura^{2,1}, Yasunobu Ogawa^{2,1}, Yoko Fukuda³, Yoshizumi Miyoshi⁴

1. 総合研究大学院大学複合科学研究科極域科学専攻、2. 国立極地研究所、3. 東京大学大学院理学系研究科地球惑星科学専攻、4. 名古屋大学宇宙地球環境研究所

1. Department of Polar Science, SOKENDAI (The Graduate University for Advanced Studies), 2. National Institute of Polar Research, 3. Department of Earth and Planetary Science, University of Tokyo, 4. Institute for Space-Earth Environmental Research, Nagoya University

The influence of the ionosphere on the aurora can be evaluated by examining the asymmetry of conjugate aurora at various time-space scales. We conducted a high-speed imaging observation of aurora at Tjornes/Iceland and Syowa/Antarctica for the time interval from 2 September 2016 to 7 September 2016 when high aurora activity continued for several days, which was driven by high-speed solar wind from a large coronal hole. It is found that an overall conjugacy is not good as we originally expected, even considering the modeled conjugate points. For example, vortex evolution from small scale (so called folds, a few ten km) to large scale (spirals, a few hundred km) occurred over whole field of view at Syowa, while such vortex structures themselves are hard to recognize at Tjornes at the same time. In this talk we discuss the conjugate morphology in more detail. Further we present the current situation and future development of the high-speed conjugate imaging observations.

キーワード：オーロラ、共役、高速撮像

Keywords: Aurora, Conjugacy, High-speed imaging