

## 原始惑星系円盤でのケイ酸塩ダストの進化

### Silicate dust evolution in protoplanetary disks

\*橘 省吾<sup>1</sup>、山本 大貴<sup>1</sup>、小林 航大<sup>1</sup>

\*Shogo Tachibana<sup>1</sup>, Daiki Yamamoto<sup>1</sup>, Kodai Kobayashi<sup>1</sup>

1. 北海道大学大学院理学研究院自然史科学専攻地球惑星システム科学分野

1. Department of Natural History Sciences, Hokkaido University

Silicate is the dominant solid component in circumstellar environments. Infrared spectroscopic observations have shown that both crystalline and amorphous silicate dust are present in protoplanetary disks, and crystalline silicate dust seems more abundant in the inner warm region of the disks. This suggests that thermal annealing of interstellar amorphous silicate dust occurred in the disk and changed the dust properties of disk dust temporally and spatially with disk evolution. Some of those processes occurred in the early Solar System may have been recorded in fine-grained matrices of less altered/metamorphosed chondrites, which contain abundant amorphous silicates and a small fraction of presolar silicate grains. Laboratory experiments help us extract the record of disk thermal processes from natural samples quantitatively. We have done experiments on crystallization and hydration experiments of amorphous silicates and evaporation and condensation experiments of crystalline silicates, focusing on kinetics of these processes. In this presentation, based on experimentally-obtained kinetic data, we will discuss the silicate dust evolution in protoplanetary disks.

キーワード：ケイ酸塩、原始惑星系円盤、速度論

Keywords: silicate, protoplanetary disk, kinetics