

Occurrence of EMIC waves and plasmaspheric plasmas derived from THEMIS observations in the outer magnetosphere

*Khan-Hyuk Kim¹, Gi-Jeong Kim¹, Dong-Hun Lee¹

1. School of Space Research, Kyung Hee University

We have statistically studied the relationship between electromagnetic ion cyclotron (EMIC) waves and cold plasmaspheric plasma (N_{sp}) in the L range of 6-12 using the Time History of Events and Macroscale Interactions during Substorms (THEMIS) data for 2008-2011. The important observational results are as follows: (1) Under quiet geomagnetic conditions ($Kp = 0-1$), the maximum occurrence rate of the hydrogen (H) band EMIC waves appears in the early morning sector (0600-0900 MLT) at the outermost region ($L = 10-12$). (2) Under moderate and disturbed conditions ($Kp > 2$), the H-band occurrence rate is higher in the morning-to-early afternoon sector for $L > 10$. (3) The high occurrence region of helium (He) band waves for $Kp = 0-1$ varies from $L = 7$ to 12 in radial distances along the local time (i.e., at $L \sim 7$ near noon and at $L = 8-12$ near late afternoon). (4) The He-band waves for $Kp > 2$ are mainly localized between 1200 and 1800 MLT with a peak around 1500-1600 MLT at $L = 8-10$. (5) N_{sp} is much higher for the He-band intervals than for the H-band intervals by a factor of 10 or more. The He-band high occurrence appears at a steep N_{sp} gradient region. (6) The morning-afternoon asymmetry of the normalized frequency seen both in H and He bands is similar to the asymmetric distribution of N_{sp} along the local time. These observations indicate that the cold plasma density plays a significant role in determining the spectral properties of EMIC waves. We discuss whether a morning-afternoon asymmetry of the EMIC wave properties can be explained by the spatial distribution of cold plasmaspheric plasma.

Keywords: EMIC waves, Plasmaspheric plasma