

Observations of dust continuum emission at 1.3mm from the protoplanetary disk around DG Tau

*百瀬 宗武¹、塚越 崇¹、スン カンロウ¹、樋口 あや²、武藤 恭之³、奥住 聡⁴、深川 美里⁵、田中 秀和⁶

*Munetake Momose¹, Takashi Tsukagoshi¹, Kang-Lou Soon¹, Aya Higuchi², Takayuki Muto³, Satoshi Okuzumi⁴, Misato Fukagawa⁵, Hidekazu Tanaka⁶

1. 茨城大学理学部、2. 理化学研究所、3. 工学院大学基礎教養・教育部門、4. 東京工業大学地球惑星科学系、5. 名古屋大学理学研究科、6. 東北大学天文学教室

1. The College of Science, Ibaraki University, 2. RIKEN, 3. Division of Liberal Arts, Kogakuin University, 4. Department of Earth and Planetary Sciences, Tokyo Institute of Technology, 5. Graduate School of Science, Nagoya University, 6. Astronomical Institute, Tohoku University

We report the ALMA long-baseline observations of dust continuum emission at 1.3mm from the protoplanetary disk around DG Tau. DG Tau is a young T Tauri star in Taurus molecular clouds. It exhibits features such as a circumstellar envelope $\sim 1000\text{au}$ in size, ionized jets, and vigorous mass accretion ($> 10^{-6} M_{\text{sun}}/\text{yr}$) onto the star. These are common to HL Tau whose protoplanetary disk was nicely imaged in the long-baseline campaign, making DG Tau an ideal object for examining the origin and universality of the ring-like features found in the HL Tau's disk. Based on the jet kinematics and gas rotation in $r < 400\text{au}$, the inclination and stellar mass were estimated to be 38° and $0.67M_{\text{sun}}$. Our present observations at the wavelength of 1.3mm have provided us with a image of dust continuum emission at the beam size of ~ 36 milli-arcseconds, or 5au . The overall structure of the disk is axisymmetric, and the surface brightness distribution is radially smooth, approximately following power-laws of $r^{-0.55}$ in $r < 12.5\text{au}$ and $r^{-1.22}$ in $12.5\text{au} < r < 50\text{au}$, respectively, with steep tapering beyond $r = 50\text{au}$. Though the inner regions ($r < 12.5\text{au}$) seems optically thick at 1.3mm, the surface density distribution in $r > 15\text{au}$ suggested from these results agrees with a classical accretion disk model. Other intriguing features are several gaps seen in $12.5\text{au} < r < 50\text{au}$, with their relative decrements of $\sim 7\%$ from the power-law distribution. These are reminiscent of spectacular concentric rings around HL Tau, and their nature should be explored further by multi-band observations with ALMA in near future.

キーワード：原始惑星系円盤、電波天文学

Keywords: protoplanetary disk, radio astronomy