

Moganite in lunar meteorite, Northwest Africa 773 clan: Trace of H₂O Ice in the Moon' s Subsurface

*鹿山 雅裕^{1,2}、富岡 尚敬³、大谷 栄治¹、瀬戸 雄介⁴、長岡 央⁵、Götze Jens⁶、三宅 亮⁷、小澤 信¹、関根 利守^{8,9}、宮原 正明⁸、留岡 和重⁴、松本 恵⁴、庄田 直起⁴、平尾 直久¹⁰

*Masahiro KAYAMA^{1,2}, Naotaka Tomioka³, Eiji Ohtani¹, Yusuke Seto⁴, Hiroshi Nagaoka⁵, Jens Götz⁶, Akira Miyake⁷, Shin Ozawa¹, Toshimori Sekine^{8,9}, Masaaki Miyahara⁸, Kazushige Tomeoka⁴, Megumi Matsumoto⁴, Naoki Shoda⁴, Naohisa Hirao¹⁰

1. 東北大学大学院理学研究科地学専攻、2. 東北大学学際科学フロンティア研究所新領域創成研究部、3. 海洋研究開発機構高知コア研究所、4. 神戸大学大学院理学研究科惑星学専攻、5. 早稲田大学先進理工学部、6. TU Bergakademie Freiberg, Institut für Mineralogie、7. 京都大学大学院理学研究科地球惑星学専攻地質学鉱物学教室、8. 広島大学大学院理学研究科地球惑星システム学専攻、9. Center for High Pressure Science and Technology Advanced Research、10. 高輝度光科学研究センター

1. Department of Earth and Planetary Material Sciences, Faculty of Science, Tohoku University, 2. Creative Interdisciplinary Research Division, Frontier Research Institute for Interdisciplinary Sciences, Tohoku University, 3. Kochi Institute for Core Sample Research, Japan Agency for Marine-Earth Science and Technology, 4. Department of Planetology, Graduate School of Science, Kobe University, 5. Research Institute for Science and Engineering, Waseda University, 6. TU Bergakademie Freiberg, Institut für Mineralogie, 7. Department of Geology and Mineralogy, Graduate School of Science, Kyoto University, 8. Department of Earth and Planetary Systems Science, Graduate School of Science, Hiroshima University, 9. Center for High Pressure Science and Technology Advanced Research, 10. Japan Synchrotron Radiation Research Institute

Lunar water locally concentrates as a result of the migration of H₂O molecules on the sunlit surface towards the colder regions. The molecular water is subsequently cold-trapped as ice on the permanently shadowed regions, the poles and theoretically Moon' s subsurface. Although a few trace of subsurface H₂O has been observed by remote sensing spectrometers (e.g., LCROSS), it has not been reported in the Apollo and Luna samples and lunar meteorites yet. In this study, lunar meteorites, the Northwest Africa (NWA) 773 clan, were investigated and thereby moganite, a monoclinic SiO₂ phase precipitated from alkaline fluids, was discovered by various microanalyses. A formation process of this lunar moganite was also interpreted to evaluate origin of the Moon' s subsurface H₂O.

Lunar meteorites of the NWA 773 clan were selected for Raman spectroscopy, electron microscopies and synchrotron X-ray diffraction (SR-XRD). The KREEP-like NWA 773 clan commonly consists of gabbroic and/or basaltic clasts.

Silica occurred as anhedral micrograins between the constituent minerals in the lunar meteorite. Raman spectra of the silica micrograins exhibited pronounced peaks at 128, 141, 217 and 503 cm⁻¹, which corresponded to those of moganite. Coesite Raman peaks were also identified together with the moganite signature. Raman intensity mapping revealed that the silica micrograins contain abundant moganite in its core, surrounded by coesite. SR-XRD of several silica micrograins also confirmed moganite and coesite. Transmission electron microscopy clarified that the silica micrograins consist of nanocrystalline particles with an average radius of 4.5 nm. Most of the SiO₂ nanoparticles were identified as moganite by selected area electron diffraction (SAED) patterns. Moganite was accompanied by small amounts of coesite, according to SAED analyses of the SiO₂ nanoparticles.

Moganite-bearing silica micrograins in the NWA 773 clan precipitated from lunar alkaline fluids rather than terrestrial weathering for the following reasons: (1) Occurrence only in a part of the NWA 773 clan. (2) Moganite surrounded by the coesite rim. (3) High moganite content contradicting reduced content to <20 wt% under dry desert condition over terrestrial age.

A formation process for lunar moganite can be explained as follows. A host gabbroic and basaltic rock of

the NWA 773 crystallised within the Procellarum KREEP Terrene (PKT). Subsequently, carbonaceous-chondrite collisions occurred on the surface of the PKT, followed by ejection of the host rock due to the impact events. The alkaline water delivered by the carbonaceous chondrite was captured as a fluid during the brecciation on the impact basin. Below the freezing point, this fluid got cold-trapped as H₂O ice in the subsurface. Simultaneously, the moganite-rich silica micrograins precipitated from the captured alkaline fluid on the sunlit surface. The NWA 773 clan was launched from the PKT by the latest impact event, thus producing transformations to coesite from moganite.

キーワード：月隕石、月、地下水、氷

Keywords: Lunar meteorite, Moon, Subsurface water, H₂O ice