

CMコンドライトの照射実験：はやぶさ2リターンサンプルの表面組織の推定

Irradiation experiments on CM chondrites: To estimate surface textures of the returned samples by Hayabusa 2

*野口 高明¹、三宅 裕二²、岡崎 隆司²、大澤 崇人³、芹澤 弘幸³、薮田 ひかる⁴、中村 智樹⁵

*Takaaki Noguchi¹, Yuji Miyake², Ryuji Okazaki², Takahito Osawa³, Hiroyuki Serizawa³, Hikaru Yabuta⁴, Tomoki Nakamura⁵

1. 九州大学基幹教育院、2. 九州大学大学院理学研究院、3. 日本原子力研究開発機構、4. 広島大学大学院理学研究科、5. 東北大学大学院理学研究科

1. Faculty of Arts and Science, Kyushu University, 2. Department of Earth and Planetary Science, Kyushu University, 3. Japan Atomic Energy Agency, 4. Department of Earth and Planetary System Science, Hiroshima University, 5. Department of Earth Sciences, Tohoku University

Introduction: In 2020, Hayabusa 2 spacecraft will return the surface and sub-surface samples from the asteroid (162173) Ryugu, a C-type asteroid. We will have an opportunity to investigate pristine materials from a C-type asteroid. Because CM chondrites contain solar gases and because most of them contain abundant subangular mineral and lithic fragments, they are regolith breccias (e.g. [1], [2], [3] and references therein). Although solar noble gases are restricted to the clastic matrix [1], [2], textures related to the solar wind irradiation and/or micrometeoroid impacts have not been identified among CM chondrites. Although there are many spectroscopic studies of CM chondrites (e.g. [4]), only a few studies are focused on the textural changes related to the micrometeoroid impacts and solar wind irradiation on CM chondrites (e.g. [5], [6]). In this study, we performed spectrum measurements, micro-petrographic study, and C K α X-ray absorption near-edge structure measurement of irradiated CM chondrites. These studies will serve to understand the space weathering on the surface of fine-grained Ryugu grains because it is highly likely that space weathering will be found on the surface of Ryugu grains.

Samples and methods: We performed irradiation of 4 keV He⁺ ions on Murchison CM chondrites at Takasaki Advanced Radiation Research Institute, Japan Atomic Energy Agency (TARRI, JAEA). The fluences are 5×10^{16} and 5×10^{17} He⁺/cm², which correspond to $\sim 10^2$ - and $\sim 10^3$ -year irradiation at 1.1 AU (the averaged orbital radius of Ryugu). Reflectance spectra of the irradiated surface were measured at JASCO Co. Ltd. by using JASCO V-670 absorption spectrometer with an integrating sphere. The irradiated samples were observed by field-emission scanning electron microscope (FE-SEM) at JAEA and Kyushu University. We observed the samples by using 2 or 3 kV acceleration voltage to avoid structural changes during observation. Thin samples were prepared by using scanning electron microscope-focused ion beam sample preparation machine and low acceleration voltage Ar milling machine at Kyushu University. They were observed by transmission electron microscope (TEM) at Kyushu University.

Results and discussion: Reflectance spectrum of the sample irradiated by a fluence of 5×10^{16} He⁺ does not show remarkable difference from the spectra of an un-irradiated sample. By contrast, a broad absorption from 0.7 to 1.4 μ m, related to the absorption by Fe-rich serpentine group minerals, is disappeared in the case of the sample irradiated with 5×10^{17} He⁺. These data suggest that 1000-year equivalent solar wind irradiation gives an effect on the shape of reflectance spectra, which is similar to the effect by dehydration [4]. There is no remarkable difference in surface morphology of the sample irradiated by a fluence of 5×10^{16} He⁺ from those of un-irradiated sample. On the other hand, the sample irradiated with 5×10^{17} He⁺ shows blistering on both matrix and chondrules. The surface of fine-grained matrix has a ~ 30 -nm thick amorphous layer. In the amorphous layer, a small amount of nanoparticles is

observed. Their 0.2-nm lattice fringes suggest that they are nanophase Fe⁰. In the case of the sample irradiated with 10¹⁷ He⁺ has ~60-nm amorphous rim containing abundant bubbles (blistering), which is especially remarkable in cronstedtite-tochilinite intergrowth. Just below the amorphous layer, both cronstedtite and tochilinite show sharp lattice fringes. The amorphous rim contains abundant nanoparticles is observed. They also show 0.2-nm lattice fringes, suggestive of nanophase Fe⁰. This result is consistent with [5].

References: [1] Nakamura T. et al. (1999a) GCA 63, 241-255. [2] Nakamura T. et al. (1999b) GCA 63, 257-273. [3] Krot A. et al. In: Meteorites and the early solar system II, pp. 679-712. [4] Hiroi T. et al. (1993) Science 261, 1016-1018. [5] Matsuoka M. et al. (2015) Icarus 254, 135-143. [5] Keller L. P. et al. (2015) LPSC 46, Abstract #1913.

キーワード：照射実験、CMコンドライト、透過電子顕微鏡

Keywords: irradiation experiment, CM chondrites, TEM